MU Guide

PUBLISHED BY UNIVERSITY OF MISSOURI EXTENSION

extension.missouri.edu

Mulches

Division of Plant Sciences

Mulches provide many benefits to plants. Benefits vary with the material used, the type of soil, the kind of plant and the cultural practices used. Mulches can also make landscapes more attractive and usable and reduce the amount of maintenance work (see Figure 1).

Why mulch?

- Surface insulation
 - to conserve moisture
 - to moderate extremes in temperature
 - to control weeds
- Soil amendment
 - to improve soil aggregation and granulation
 - to increase water absorption and retention
 - to prevent soil compaction and improve aeration
- Beautification
 - to make surface areas more attractive
 - to make surface areas more usable for paths, play and sitting areas
 - to make areas easier to maintain

Types of mulch

Many types of material, organic and inorganic, may be used as mulch. Organic mulches may break down in one season or less or persist for more than one season. The first type is used mainly in vegetable and flower gardens or around newly planted trees and shrubs. They are usually decomposed enough by the end of the growing season that they can be spaded or plowed under, increasing the organic matter content of the soil and thereby improving soil structure. Even if not incorporated, mulch improves the soil as worms and soil microbes move it into the soil. Some common mulches of this type are listed in Table 1.

Organic mulches that persist for more than one growing season are more useful around permanent plants. These are summarized in Table 2.

Author

Christopher J. Starbuck, Associate Professor, Division of Plant Sciences, is a State Horticulture Extension Specialist.







Figure 1. Mulch beds can provide a unifying effect in the landscape (top) while keeping the soil cooler in summer and warmer in winter, an important benefit for flowering bulbs (middle). Mulch can also make an attractive walkway through the landscape (bottom).



Figure 2. One use for an inorganic mulch, such as gravel, is in a rock garden.

Inorganic mulches may be gravel, crushed stone or some manufactured product (Figure 2). Most of these materials are not as beneficial to plants as the organic mulches are because they do not improve the structure or nutrient content of the soil. Black plastic mulch is not recommended for use in the landscape but may be useful in the vegetable garden. Spun-bonded or woven landscape fabrics (geotextiles) are better suited for landscape purposes. Normally these fabrics are cov-

Table 1. Relative value of mulches that break down in one season or less.

Material	Resistance to compaction	Attractiveness	Resistance to wind blowing	Availability	Source of weeds and disease	Fire hazard	Comments
Compost	В	В	Α	Α	С	No	Value varies with ingredients.
Corn stalks	Α	F	В	С	Α	Yes	Coarse and unsightly. Should shred or compost.
Hay	В	С	В	С	С	Yes	Good when available.
Lawn clippings	D	D	В	Α	С	No	Compost first.
Leaf mold	В	В	Α	Α	С	No	Value varies with ingredients.
Leaves	F	В	D	Α	В	Yes	Compost first.
Manure (well rotted)	В	В	Α	С	F	No	Odor may be bad.
Peat moss	В	Α	Α	Α	В	No	Universally available. Coarse grades best.
Rice hulls	В	С	D	D	С	No	Good when available.
Straw	Α	D	D	В	D	Yes	Often contains grain seed.
Waste paper	С	F	F	Α	Α	Yes	Must be shredded. Unsightly.
Key: A = Excellent; B = Good; C = Fair; D = Poor;							

F = Unsatisfactory

ered with a more decorative mulch rather than used alone. However, soil structure may deteriorate under geotextiles, because the organic mulch above it cannot enter the soil. Some commonly used inorganic mulches are summarized in Table 3.

Though not generally considered mulches, some low-growing groundcover plants produce many of the same benefits as mulch. They shade the soil surface, keeping it cool in summer and at the same time preventing evaporation. For more information on groundcovers, see MU publication G6835, Selected Groundcovers for Missouri.

Using mulches

On the whole, mulches should not be considered a fertilizer. Most of them release some nutrients as they decompose, but the fertilizer value is very small in relation to the physical effects. Where mulches are needed, use them for the mulch value and add fertilizer as a supplement.

The effectiveness of a mulch depends on the density and texture of the material used and the depth of the layer applied. A 2- to 4-inch layer of most products is needed to insulate the soil effectively and prevent weed growth.

Table 2. Relative value of some persistent mulches.

	Resistance to compaction	eness	Resistance to wind blowing	lity	Source of weeds and disease	ard	
Material	Resista	Attractiveness	Resista	Availability	Source	Fire hazard	Comments
Bark chunks	Α	Α	Α	Α	Α	No	Generally available. Expensive.
Cocoa shells	Α	Α	В	D	Α	No	High potassium content may cause problems.
Corn cobs	В	D	В	С	В	No	Add nitrogen. Unsightly.
Cottonseed hulls	В	С	D	D	В	No	Hard to keep in place.
Pecan hulls	Α	Α	В	С	Α	No	Locally available. Good.
Pine needles	Α	В	В	С	В	Yes	Especially good on acid- loving plants.
Sawdust (coarse)	С	С	В	В	Α	No	Add nitrogen. Use aged material.
Sawdust (fine)	D	С	D	В	Α	No	Compacts too easily. Requires nitrogen.
Shredded bark	В	Α	Α	В	Α	No	Good when available.
Wood chips	В	В	Α	В	Α	No	Inexpensive when available.
Wood shavings	В	В	В	С	Α	Yes	Add nitrogen when spaded under.
Wood fibers	С	O	В	D	Α	No	Add nitrogen. Coarse grade preferred.
Key: A = Excellent; B = Good; C = Fair; D = Poor;							

F = Unsatisfactory

Table 3. Inorganic mulching materials.

Material	Resistance to compaction	Attractiveness	Resistance to wind blowing	Availability	Source of weeds and disease	Fire hazard	Comments
Black plastic film	Tears	F	F	Α	Α	No	Must anchor. Unsightly.
Crushed rock	A	В	Α	Α	Α	No	Many colors available. Avoid crushed limestone
Geotextile weed barrier	А	F	F	Α	Α	No	Must anchor. Best covered by other mulch.
Gravel	Α	В	Α	Α	Α	No	Use sparingly.
Volcanic rock	Α	В	Α	В	Α	No	Small sizes moved by water.
Perlite	В	В	D	Α	Α	No	Good as soil amendment.
Vermiculite	С	С	D	Α	Α	No	Physical structure breakdown.
Key: A = Excellent; B = Good; C = Fair; D = Poor; F = Unsatisfactory							

Mulching suggestions

Considering the growth requirements of the multitude of garden and landscape plants, obviously there is no one mulch that will be superior in all cases. The following general suggestions should help in selecting the right mulch.

Bulbs

Among garden bulbs, lilies benefit more from mulching than any other group. Two to 3 inches of wood shavings, straw, chopped hay or other lightweight material serve to lower the temperature of the soil in summer, conserve moisture and give winter protection to the more tender species.

Most other bulbs are sufficiently winter hardy but benefit from mulching for other reasons. Small, shallow-planted bulbs can be heaved out of the soil by alternate freezing and thawing. This effect can be largely prevented by light mulching. Mulches generally improve the appearance of flowering bulbs growing in a bed of otherwise barren soil. They offer a background for viewing the new spring growth and also reduce the spattering of plants with mud during rainstorms.

Evergreens

Plants that most need mulching are evergreen shrubs and young evergreen trees, especially the broad-leaved types. Mulches help prevent winter drying of these plants. Winter drying causes foliage to become scorched and discolored over winter and may cause the plant to die. Drying develops when foliage loses moisture faster than it can be taken up by the roots from a dry or frozen soil.

Plants that should almost always be mulched are azalea, boxwood, Chinese and Japanese hollies, Japanese andromeda and rhododendron. Exotic evergreens and those planted north of their best zone of adaptation should be kept well mulched year-round.

Coarse mulches should be used: wood chips, bark products, shavings, chopped corn cobs, pine needles, hedge trimmings or partially finished coarse compost. Mulches should be renewed in early winter. A layer 2 to 4 inches deep is adequate in most cases but can be more or less, depending upon site, soil, climatic zone and the plant itself. The mulched area should be at least as broad as the plant itself and preferably broader since most woody plants have root systems that reach beyond the spread of the branches.

Fruits

To keep young fruit trees growing well, maintain a mulched area 3 to 4 feet in radius around the base. Coarse materials such as hay, straw, wood or bark products are most useful. Keep mulch pulled away from the trunk of the tree. The mulch can be replenished in late March or early April before new weed growth appears. Grapes and berry bushes may be kept permanently mulched.

Strawberries should be mulched preferably with clean straw or with prairie hay in mid-December. Use enough mulch, loosely scattered, to barely hide the foliage. In spring when new leaves begin to grow, remove enough of the mulch to allow the plants to emerge.

Perennials

Most garden perennials benefit from summer mulches for preserving soil moisture and for reducing soil temperature. Perennials that most often need winter protection in Missouri include baby's breath, bellflower, chrysanthemums, delphinium, hardy gloxinia, hypericum, linum, lupine, oriental poppy, Stoke's aster, thrift and thyme.

Some perennials that are normally cold hardy are often damaged by alternate freezing and thawing of the soil, which may loosen or heave the plants and expose them to drying. Among these are aconite, coral bells and newly planted peony and plantain lily.

A few plants are extremely sensitive to damage by high summer soil temperatures or drought. Delphinium, lupine, primrose and Siberian iris are among the most troublesome in this respect.

Two to 3 inches of straw, hay, wood shavings or partially composted leaves would be an appropriate mulch for most perennials. Pine needles or evergreen boughs are also effective. When using a heavy layer of mulch on perennials, gradually remove it during the onset of warmer weather in late winter. Heavy mulch left too long may cause early emergence of weak, ten-

der growth. Such growth may be killed or permanently injured by late freezes.

Roses

Experienced rose growers recognize the improvement in vigor and appearance of roses as a result of summer mulching. The mulch layer should be about 2 inches deep and of fairly coarse material such as unfinished compost, aged sawdust, bark, milled wood fiber or tobacco stems.

By mid-November, provide adequate winter protection with a mound of finished compost, garden soil or coarse peat. The mound should be about 8 inches deep across the center of the plant. Mounds should be removed in early spring when overnight temperatures below about 28 degrees F are no longer expected. If later, unseasonably cold weather threatens and new, tender shoots have begun to grow, hasty covering may be needed.

Trees and shrubs

Trees and shrubs that gain most from mulching are new plantings, ones with shallow roots, and ones that suffer from extended drought or are otherwise damaged by high soil temperatures (Figure 3). Flowering dogwood and sugar maple are examples of plants whose leaves scorch easily in dry weather. Some leaves of birches, willow and tulip trees often yellow and drop during extended drought.

All trees and shrubs should be mulched at planting time to aid them in becoming established. A mulched area around the base of trees and shrubs makes mowing easier and helps prevent mower damage to the plant.

The material used for mulching is perhaps less important than the simple fact that the plants are mulched. Most coarse materials are satisfactory. Avoid materials such as sawdust, which packs tightly, and those too unsightly in the landscape. Planting groups of trees and shrubs together in mulched beds is efficient and simulates natural conditions for most species (Figure 4).

Vegetables and annual flower gardens

The primary reason for mulching vegetable and flower gardens is to conserve soil moisture. Crops most practical to mulch are beans, peppers, sweet corn, tomatoes, vine crops and other long-term summer crops with large amounts of foliage. Long-term crops started in the cool season, including broccoli, cabbage, cauliflower and potatoes, will also yield better when



Figure 3. Newly planted trees benefit greatly from a ring of mulch around the base of the tree.



Figure 4. Trees and shrubs can be grouped in mulched beds.

they are mulched. Asparagus and rhubarb should be kept constantly mulched to conserve soil moisture and reduce weed problems.

Some of the best mulches for annual vegetable crops are composts, straw, hay or other materials that will be largely decomposed at the end of the season.

UNIVERSITY OF MISSOURI
susued in furtherance of Cooperative Extension Work Acts of May 8 and June 30, 1914, in cooperation with the United States Department of Agriculture. Michael D. Ouart, Director, Cooperative Extension, University of Missouri, Columbia, MO 65211. ■ University of Missouri Extension does not discriminate on the basis of race, color, national origin, sex, sexual orientation, religion, age, disability or status as a Vietnam era veteran in employment or programs. If you have special needs as addressed by the Americans with Disabilities Act and need Vietnam era veteran in employment or programs. If you have special needs as addressed by the Americans with Disabilities Act and need Vietnam era veteran in employment or programs. If you have special needs as addressed by the Americans with Disabilities Act and need Vietnam era veteran in employment or programs. this publication in an alternative format, write ADA Officer, Extension and Agricultural Information, 1-98 Agriculture Building, Columbia, MO 65211, or call 573-882-7216. Reasonable efforts will be made to accommodate your special needs.